## Killing symmetries of generalized Minkowski spaces. 1- Algebraic-infinitesimal structure of space-time rotation groups

Fabio Cardone<sup>a,b</sup>, Alessio Marrani<sup>c,d</sup> and Roberto Mignani<sup>b-d</sup>

> a Dipartimento di Fisica Università dell'Aquila Via Vetoio
> 67010 Coppito, L'Aquila, Italy
> b I.N.D.A.M. - G.N.F.M.
> c Università degli Studi "Roma Tre" Via della Vasca Navale, 84 00146 ROMA, Italy
> d I.N.F.N. - Sezione di Roma III

> > July 30, 2013

## Abstract

In this paper, we introduce the concept of N-dimensional generalized Minkowski space, i.e. a space endowed with a (in general nondiagonal) metric tensor, whose coefficients do depend on a set of nonmetrical coordinates. This is the first of a series of papers devoted to the investigation of the Killing symmetries of generalized Minkowski spaces. In particular, we discuss here the infinitesimal-algebraic structure of the space-time rotations in such spaces. It is shown that the maximal Killing group of these spaces is the direct product of a generalized Lorentz group and a generalized translation group. We derive the explicit form of the generators of the generalized Lorentz group in the self-representation and their related, generalized Lorentz algebra. The results obtained are specialized to the case of a 4-dimensional, "deformed" Minkowski space  $\widetilde{M}_4$ , i.e. a pseudoeuclidean space with metric coefficients depending on energy.

## 1 Introduction

In the last years, two of the present authors (F.C. and R.M.) proposed a generalization of *Standard Special Relativity* (SR) based on a "deformation"

of space-time, assumed to be endowed with a metric whose coefficients depend on the energy of the process considered [1]. Such a formalism (*Deformed Special Relativity*, DSR) applies in principle to *all* four interactions (electromagnetic, weak, strong and gravitational) - at least as far as their nonlocal behavior and nonpotential part is concerned - and provides a metric representation of them (at least for the process and in the energy range considered) ([1]-[5]). Moreover, it was shown that such a formalism is actu-

ally a five-dimensional one, in the sense that the deformed Minkowski space

is embedded in a larger Riemannian manifold, with energy as fifth dimension [6].

In this paper, following the line of mathematical-formal research started with [7] and [8], we introduce the concept of N-dimensional generalized Minkowski space, i.e. a space endowed with a (in general non-diagonal) metric tensor, whose coefficients do depend on a set of non-metrical coodinates. The deformed space-time  $\widetilde{M}_4$  of DSR is just a special case of such